



Tiny teeth!

A short guide to healthy teeth for the under fives

Toothy tips for tiny tots


A happy child with a bright smile is what every parent wants to see.

Good teeth help your child feel confident but tooth decay can cause pain and worry, even in small children.

When it comes to children's teeth, regular brushing and visits to the dentist are important.

What, when and how your child eats and drinks is also important and this booklet gives simple facts on how you can keep your little ones grinning from ear to ear.

It's a toddler!



First steps

Tooth decay (also known as dental caries) is caused when the bacteria in our mouth turn sugars left behind after eating and drinking into acid. This causes damage to the teeth, leading to decay.

You can help protect small children's teeth in 3 simple steps

1. Keep sugary foods and drinks to mealtimes
2. Brush teeth and gums twice a day for at least 2 minutes with a fluoride toothpaste
3. Register with a dentist and visit regularly; the sooner the better! Children are treated free under the NHS

Reducing sugar in your child's diet will help **prevent tooth decay**; it's how often that it's served which is particularly important.

The more often we eat and drink sugary things the more acid is produced and the more damage is caused to tiny teeth. Sweet drinks in a bottle, particularly at bedtime, or sweets like lollipops will damage teeth because the sugar will be in contact with the teeth for a long time.



Erosion explosion

Sugar is not the only issue for dental health. Acidic drinks can also be a big problem for tiny teeth by wearing away the surface of the tooth.

Fizzy drinks, fruit juices and squashes are often the acid culprits in the erosion explosion, even the sugar-free or diet varieties.

To help avoid tooth erosion, don't offer children too many acidic drinks and keep them to mealtimes rather than between meals.

Only offer water or milk (breastmilk or formula to infants) between meals – these are good choices for tiny teeth.


It is best to give tiny tots water or milk to drink at mealtimes too. Fruit juices are acidic and contain sugar and are not recommended. If you do give them, make sure they are well diluted with water and they are in a cup, not a bottle or lidded cup with a valve. This makes it less likely to cause tooth erosion.

Fizzy drinks should be avoided. Squashes and other soft drinks are not recommended for babies and toddlers. If you choose to give these to your child make sure they are:

- well-diluted
- drunk from a cup
- drunk at mealtimes
- never given in a bottle or at bedtime.

If they are thirsty at bedtime or during the night offer water, and avoid giving your child juices or sugary foods or drinks.

Remember, the more often tots eat sugary foods or have sugary or acidic drinks the more damage is caused to tiny teeth.



Toothpaste power!

Take care of tiny teeth



For a sweet smile, brush teeth twice a day for at least 2 minutes with fluoride toothpaste* and discourage a sweet tooth.

Start to brush your baby's teeth as soon as they begin to come through. Parents should continue to brush tiny teeth or supervise brushing until at least seven years of age.


* see page 18

Say cheese

Milk provides calcium which is needed for strong teeth.

Try swapping biscuits and cakes for tooth-friendly snacks like small cubes or sticks of hard cheese and plain yogurt – like milk they also have calcium.

Sweet snacks mean tooth decay attacks!



Look out for secret sugar

Sugar comes in many different disguises. It can be labelled in a different way so as well as sugar, look out for sucrose, fructose, glucose, maltose, dextrose, honey and syrups, and concentrated fruit juice.

Smiles of snack ideas



- Peeled and chopped fruit and vegetables
- Plain yogurt with added fruit
- Crumpets, toast fingers and cheese spread
- Homemade plain popcorn
- Chapatti, pitta bread, potato cakes, matzos and rice cakes
- Plain biscuits like water biscuits and cream crackers
- A stick of cheese

Bites for growing mites

Milky goodness



As well as providing calcium which is needed for strong teeth and bones, milk provides much more.

- Milk is packed with nutrients providing:
- Iodine for children's growth
- Phosphorus for energy release
- Protein for bone development
- Potassium for nerve function
- Vitamin B2 for healthy skin and vision
- Vitamin B12 for red blood cells.

From weaning you can use whole cow's milk in cooking or for mixing into cereal, and you can also give foods made from whole milk like yogurt, fromage frais and cheese. From 12 months whole cow's milk is suitable as the main drink, but not before this time.

From age 2 years onwards you can start introducing semi-skimmed milk if your child is eating a variety of foods and growing well. Otherwise stick to whole milk. 1% fat or skimmed milk are not suitable for under 5s.



Family recipes

with milky, cheesy or yogurty ingredients

Easy Peasy Bolognese

- 1 tablespoon olive oil**
- 1 onion, peeled and diced**
- 1 garlic clove, crushed**
- 1 carrot, peeled and diced**
- 1 pepper, deseeded and diced**
- 1 teaspoon dried mixed herbs**
- 400g lean minced beef**
- 120ml milk**
- 2 x 400g chopped tomatoes**
- 300g mini pasta shapes**
- 15g freshly grated Cheddar or Parmesan, to serve**

Heat the oil in a large saucepan over medium heat. Add the onion, garlic, carrot, pepper and dried herbs, and fry for about 8 minutes, until softened. Add the minced beef and cook until browned, stirring to break it up. Add the milk and tomatoes and bring to the boil. Turn down the heat and simmer with the lid on for 30 minutes. If the sauce is still too juicy take the lid off and cook for longer.

Cook the pasta according to packet instructions and serve with the Bolognese sauce and some Cheddar or Parmesan cheese grated on top.

Easy, cheesy tuna and potato bake

350g potatoes, washed and cut into similar sized pieces

25g butter

25g flour

1 can of tuna (185g), drained

285ml (½ pint) milk

1 large leek, washed and chopped

1 red pepper, washed and chopped

90g Lancashire or other white cheese

Preheat the oven to 200°C or gas mark 6. Parboil the potatoes leaving the skins on, and cut into thin wedges. Melt the butter in a pan; add the chopped leeks and pepper, and cook until just soft. Stir in the flour, cook for 1 minute, add the milk gradually, whisk continuously and heat gently until the sauce begins to thicken. Add the tuna and turn into an ovenproof dish. Arrange the potato wedges on the top and grate or crumble the cheese over. Bake for 20 minutes until the cheese is golden brown.

Cheese & pineapple sticks

15-20g of cheese

1 large slice of fresh pineapple - cut into pieces

Cocktail sticks


Add 2 pieces of pineapple to the stick and 1 piece of cheese, add another 2 pieces of pineapple. Repeat until ingredients are finished.

Yogurt and strawberries

80-100g of natural unsweetened yogurt

7 fresh strawberries chopped

Place yogurt in a bowl and add chopped strawberries.



Serves 2

Banilla whizz

- 1 large ripe banana, sliced**
- 1 small pot (approx 150g)
thick and creamy vanilla yogurt**
- 450ml (¾ pint) milk**

Reserve two pieces of banana for decoration, then put the remainder into a blender or food processor with the yogurt and milk. Blend for 10-15 seconds until smooth.

Pour into two glasses and decorate with the reserved banana.


Tasty tip

If you can't find vanilla yogurt, use natural yogurt and add 1-2 drops of vanilla extract to the blender with the other ingredients.

Serves 4

Milky wobble

- 1 pack of sugar-free jelly (135g)**
- 125ml boiling water**
- 250ml whole milk**



Dissolve the jelly in boiling water. Once cooled but not set, whisk in the milk using a fork. Pour into a mould or container and leave to set in a fridge.

Your questions answered

My child loves dried fruit. Is it OK to give this as a snack?

Dried fruit is nutritious but should be eaten with a meal and not as a snack. The natural sugars in dried fruit can cause tooth decay. Eating dried fruit at mealtimes makes damage to tiny teeth less likely.



I've heard bottles are bad for teeth. When should I stop giving my baby a bottle?

It's fine to use a bottle for milk feeds up until your baby is around 12 months old, however, prolonged bottle feeding is not recommended because older babies and toddlers can begin to rely on bottle drinking for comfort. Sweet drinks in bottles are very damaging to teeth. It's best not to put anything in a bottle other than expressed breastmilk, formula or water. As soon as your child is ready, encourage them to use an open cup.

Should I stop giving my child fruit juice?

New guidelines do not recommend fruit juice at any age because its sugar and acid content is harmful to teeth. If you give your child fruit juice ensure it is well diluted and taken at mealtimes only.

I think my child's allergic to cow's milk. Should I change to a soya drink?

Allergy to cow's milk is not as common as people think, affecting about 1 in 50 infants, but it is much less common in older children (most have outgrown their allergy by the time they start school). Always get advice from a Registered Dietitian (through your GP) before excluding cow's milk.

Soya products should only be used if advised by a dietitian or GP, as children who are allergic to cow's milk may also be allergic to these. If you do give soya-based drinks it is important to choose one with added calcium, and remember that some soya-based drinks contain added sugars, which are not good for teeth.

Are sugar-free squashes safe to give my toddler between meals?

Diet, reduced-sugar or no added sugar drinks can encourage a sweet tooth and usually contain artificial sweeteners which are not recommended for toddlers. Your toddler should avoid all soft drinks such as fizzy drinks and squashes even if sugar-free. These drinks can still contribute to tooth decay and provide little nutritional value.

Is it OK to give my child tea or coffee?

Giving tea or coffee to young children reduces their absorption of iron and is not recommended. They also contain caffeine which is a stimulant and can disturb children's sleep.

My child keeps asking for sweets and fizzy drinks! What can I do?

Sweets and fizzy drinks can harm your child's teeth, as well as taking away their appetite for more nutritious foods. It's not advisable to give fizzy drinks to toddlers. It's OK to give your toddler sweets occasionally, but try not to offer them regularly so that your child won't expect them and you'll avoid tantrums. If you do let your child have sweets it's best to give them all at once, at the end of a meal, as this is less harmful for their tiny teeth. Also have plenty of alternative snacks available.

What type of toothpaste is best for young children?

For children under three years you can use a smear of family toothpaste containing at least 1,000ppm (parts per million) fluoride; for children aged three to six years a pea-sized blob is needed containing 1,350-1,500ppm fluoride. Check the toothpaste packet for this information or ask your dentist. Always supervise toothpaste use and help your child brush their teeth until at least seven years old. Also make sure they don't eat toothpaste from the tube! Your dentist can also give you advice on whether your child would benefit from extra protection such as fluoride varnishes.

My child makes it difficult for me to brush his teeth. What can I do?

When it comes to tooth brushing, it is best to start young – as soon as your baby's teeth appear. Make it a fun part of your child's morning and bedtime routines – sing songs while you brush and use different coloured or cartoon-themed toothbrushes.

References

- NHS Choices. Looking after your infant's teeth
<http://www.nhs.uk/Conditions/pregnancy-and-baby/Pages/looking-after-your-infants-teeth.aspx>
[accessed 08/2016]
- NHS Choices. Kids' teeth Q & As
<http://www.nhs.uk/Livewell/dentalhealth/Pages/Goodhabitskids.aspx>
[accessed 08/2016]
- NHS Choices. Children's Teeth
<http://www.nhs.uk/Livewell/dentalhealth/Pages/Careofkidsteeth.aspx>
[accessed 08/2016]
- NHS Choices. Drinks and cups for children
<http://www.nhs.uk/Conditions/pregnancy-and-baby/Pages/drinks-and-cups-children.aspx>
[accessed 08/2016]
- NHS Scotland. Childsmile Parents and Carers
<http://www.child-smile.org.uk/parents-and-carers/index.aspx>
[accessed 08/2016]
- The Department of Health (2014) Delivering better oral health: an evidence-based toolkit for prevention, Third Edition
https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/367563/DBOHv32014OCTMainDocument_3.pdf
[accessed 08/2016]
- British Dental Health Foundation. Caring for teeth: Diet
<http://www.dentalhealth.org/tell-me-about/topic/caring-for-teeth/diet>
[accessed 08/2016]
- BUPA. Caring for your child's teeth (last reviewed: 04/2013)
<http://www.bupa.co.uk/individuals/health-information/directory/d/child-dental-care>
[accessed 08/2016]

For details on additional information sources please contact The Dairy Council

Last reviewed: 01/2017
Next review due: 01/2018



Tel 020 7025 0569
info@dairycouncil.org.uk

For **free** copies of The Dairy Council's publications visit www.milk.co.uk